

Original Research

Evaluation of Risk Factors for Fever & Convulsion in Pediatrics 6 Months to 6 Years: A Review Study

Jalal Nourmohammadi¹, Maryam Khalili², Dorrin Nikbakht^{3*}

1. Master of Nursing, Pediatrics Department, Shahid Hasheminejad Medical Research Center, Mashhad University of Medical Sciences, Iran
2. Master of Nursing, Intensive Care Department, Iran University of Medical Sciences, Iran
3. Instructor of Operating Room, Department of Anesthesiology, School of Allied Medical Sciences, Alborz University of Medical Sciences, Iran

***Corresponding Author: Dorrin Nikbakht**, Instructor of Operating Room, Department of Anesthesiology, School of Allied Medical Sciences, Alborz University of Medical Sciences, Iran. E-mail: dorrinblue@gmail.com. Orcid:0000-0002-2902-1927

Abstract:

Background:

Fever & Convulsion are the most common type of seizures and neurological problems of childhood and one of the most common reasons for admission to the emergency department and Neurology is a children's hospital. The aim of this study was to investigate the risk factors for Fever & Convulsion in children aged 6 months to 6 years.

Methods:

In total, 43 articles with a time limit of the last 5 years were obtained by deleting 9 articles whose full text not available; finally, 34 articles were included in the study

Results:

Risk factors for Fever & Convulsion in children 6 months to 6 years old using a review of studies conducted in the period 2018 to 2022. Findings of this study showed that genetics, gender, iron deficiency anemia, previous history of Fever & Convulsion, young age (under one year), infections and perinatal complications are effective in the development of fever and seizures and are associated with hospitalization of the child.

Conclusion:

The results showed that various factors are effective in the development of febrile seizures in children aged 6 months to 6 years. Fortunately, Fever & Convulsion are a benign disease in children that does not have long-term neurological complications and has a good prognosis.

Keywords: Fever & Convulsion, Pediatrics, Risk Factor

Submitted: 14 August 2022, Revised: 27 September 2022, Accepted: 27 October 2022

Introduction

Fever & Convulsion are the most common type of seizures and neurological problems of childhood and one of the most common reasons for admission to the emergency department and Neurology is a children's hospital (1). By definition, the disease is an event that occurs in infancy and early childhood, usually between 6 months and 6 years of age at a temperature of more than 38.5 ° C, and evidence of central nervous system infection or water and electrolyte disturbances and history no previous seizures without fever (2).

Fever & Convulsion can occur in any form, but most cases are generalized tonic-clonic seizures that resolve spontaneously after a few minutes (3). The age of onset of this disorder is 14 to 18 months and is seen in 75% of cases under the age of 3 years (4). The prevalence of Fever & Convulsion in different societies is reported to be between 2 and 14%, for example in Europe and the United States, 2 to 5%, India 5-10%, in Japan 8.8% and in Ghana 14% (5).

Although this event is benign and has a good prognosis, fever is one of the most important public health concerns that occurs frequently during childhood and children with Fever & Convulsion develop more febrile episodes than children. Peers experience no Fever & Convulsion (6). Most Fever & Convulsion are seen in normal children, although children with neurological or developmental disorders may be more susceptible (7).

This type of seizure is usually seen in the first hours of infectious disease and its maximum incidence (73%) is in the first 24 hours of fever (8). The onset of seizures one day after the onset of febrile illness suggests another possible diagnosis for the child, such as meningitis and electrolyte disturbance (9).

The disease originates from environmental and genetic factors, although the main cause of this type of seizure is still unknown (10). But factors such as a positive family history, perinatal events, developmental delay, male

gender, duration of breastfeeding, high temperature, low birth weight, elevated blood bilirubin, or infants whose mothers use alcohol and cigarettes; Has been reported in febrile seizures (11).

The prognosis for Fever & Convulsion is very good, and long-term antiepileptic drugs are not needed for simple Fever & Convulsion and in most cases for complex Fever & Convulsion. The aim of this study was to get acquainted with the risk factors for Fever & Convulsion in children and to improve child care.

Methods

This review study by searching the databases of Google Scholar, Scopus, PubMed and Web of Science, using Persian keywords Fever & Convulsion, Pediatrics, Risk Factor Their English was done. In total, 43 articles with a time limit of the last 5 years were obtained by deleting 9 articles whose full text not available; finally 34 articles were included in the study

Results

The aim of this study was to determine the risk factors for Fever & Convulsion in children 6 months to 6 years old using a review of studies conducted in the period 2018 to 2022. Findings of this study showed that genetics, gender, iron deficiency anemia, previous history of fever and seizures, young age (under one year), infections and perinatal complications are effective in the development of fever and seizures and are associated with hospitalization of the child.

In the study by Gvozdenovic et al., The body temperature was 39.7 ° C during seizures caused by moderate fever. According to studies, certain genetic backgrounds are involved in the development of febrile seizures. There was a significant relationship between the presence of seizures in the first degree family and its recurrence. This association has been reported in studies (12). This association

has been reported in studies by Adebukola et al, Alfahid et al, And Kaboré et al (13) (14) (15). Habeeb concluded in his study that children with Fever & Convulsion are more likely to have a seizure if they have a positive family history of Fever & Convulsion (16). A positive family history was associated with Fever & Convulsion in 25 to 40% of cases, and a prevalence of 9 to 22% of reported siblings was reported (17).

Fever & Convulsion were significantly higher in boys than in girls. 60% of the cases were boys and 40% of the cases were girls (18). In the study of Alruwaili et al., The prevalence of fever and seizures was higher in boys (19). In the study by Alqudah et al., 54% of patients had seizures due to male fever (20).

Fever & Convulsion at a age of less than one year are a risk factor. Because children who have seizures at a younger age have a longer duration in the age group of people who are at risk for Fever & Convulsion are located (21). In the study of Prakash et al., The highest frequency of Fever & Convulsion was observed in children under two years of age (22).

A history of perinatal problems was associated with Fever & Convulsion in children (23). In the study Laino et al, Perinatal problems such as preterm delivery and low birth weight were identified as risk factors for Fever & Convulsion (24). In the study by Mosili et al, the role of developmental delay as a risk factor in the development of Fever & Convulsion was proven (25).

The results of studies indicate that plasma ferritin levels were significantly lower in children with Fever & Convulsion. This may indicate the possible role of iron deficiency in its occurrence. The prevalence of Fever & Convulsion is very low in children with thalassemia major with high iron levels (26) (27).

In Hameed et al study on children with Fever & Convulsion, serum zinc levels in the patient

group were significantly lower than serum zinc levels in the control group. There was also a difference in serum zinc levels in the cerebrospinal fluid of both groups (28) (29). In the study of Hawas et al, It was performed on children with seizures and non-children. The cerebrospinal fluid of both groups was examined for magnesium and calcium levels and the result was that the level of magnesium in the cerebrospinal fluid of the patient group was far from there was less evidence (30).

The increase in the number of viral infections in children has been considered as a cause of Fever & Convulsion in the last decade. In Asian studies, influenza Type A virus has been the most common cause of seizures. Herpesvirus Type 6 infection has been the most common cause of Fever & Convulsion in children in the United States (31).

The relative risk of Fever & Convulsion temporarily increases within 2 weeks after measles and mumps (MMR) vaccination. The rate of the first Fever & Convulsion in children who receive the vaccine is 10% higher than others (32). There is also a risk of 8 to 14 days after MMR vaccination. Fever & Convulsion increase (33). While this increased risk for diphtheria-tetanus-pertussis vaccination occurs on the day of vaccination (34).

Conclusions

The results showed that various factors are effective in the incidence of Fever & Convulsion in children aged 6 months to 6 years. Fortunately, Fever & Convulsion are a benign disease in children that does not have long-term neurological complications and has a good prognosis. The results of this study can be used to educate nurses, nursing parents and parents to identify risk factors for febrile seizures and Standard nursing care used during pediatric hospitalization for nurses working in pediatric wards.

Declaration of Competing Interest:

The authors report no declarations of interest

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Ethics statement

Written informed consent was obtained from the individuals for the publication of any potentially identifiable data included in this article.

References

1. Duthie, L. and R. Begley, Febrile convulsions. *Paediatrics and Child Health*, 2021. 31(11): p. 415-418.
2. Al Jandale, O., et al., Tonic non-colonic convulsions-status epilepticus- as the presenting complaint of COVID-19. *Annals of Medicine and Surgery*, 2022. 78: p. 103744.
3. Canpolat, M., et al., Investigating the prevalence of febrile convulsion in Kayseri, Turkey: An assessment of the risk factors for recurrence of febrile convulsion and for development of epilepsy. *Seizure*, 2018. 55: p. 36-47.
4. Yonemoto, K., K. Okanari, and H. Koga, Optimal Doses of H1 Antihistamines Do Not Increase Susceptibility to Febrile Convulsions in Children. *Pediatric Neurology*, 2018. 87: p. 42-47.
5. Shi, K., et al., Risk factors for the recurrence of convulsions with mild gastroenteritis in children. *Seizure*, 2020. 80: p. 192-195.
6. Green, R., et al., Management of acute fever in children: Consensus recommendations for community and primary healthcare providers in sub-Saharan Africa. *African Journal of Emergency Medicine*, 2021. 11(2): p. 283-296.
7. Jiang, L., et al., Reversible splenic lesion syndrome in children with benign convulsions associated with mild gastroenteritis: A retrospective study of five cases. *Brain and Development*, 2019. 41(3): p. 271-275.
8. Osman, N.M. and J.P.G. Gai, Assessment of the necessity of routine lumbar puncture among children with fever and convulsions. *Sudan Journal of Medical Sciences*, 2019. 14(3): p. 162-171.
9. Hao, X.-s., et al., Clinical characteristics, treatment, and long-term outcomes in children suffering from benign convulsions with mild gastroenteritis: a retrospective study. *BMC pediatrics*, 2020. 20(1): p. 1-6.
10. Orji, M.-L., et al., Prevalence, pattern and outcome of pediatric Lassa fever disease (LFD) in a tertiary hospital, southeast Nigeria. 2020.
11. Kavanagh, F.A., et al., Recognition and management of febrile convulsions in children. *British Journal of Nursing*, 2018. 27(20): p. 1156-1162.
12. Gvozdenovic, E., et al., Impact of history of febrile convulsions on the risk difference of febrile convulsions with the tetravalent measles-mumps-rubella-varicella vaccine: Post-hoc exploratory analysis of results from a matched-cohort study. *Vaccine*, 2018. 36(39): p. 5803-5806.
13. Adebukola, O.O., et al., Pre-hospital management of childhood convulsions in Ilesa, South-west, Nigeria. *African Journal of Medical and Health Sciences*, 2021. 20(1): p. 1-7.
14. Alfahid, F., et al., Knowledge, attitude, and practice of parents toward febrile convulsions in Saudi Arabia, 2019. *Age*, 2020. 30: p. 30-45.
15. Kaboré, A., et al., Febrile Convulsions in Infants at the Pediatrics University Hospital Center Charles de Gaulle of Ouagadougou (Burkina Faso). *Open Journal of Pediatrics*, 2018. 8(2): p. 199-206.

16. Habeeb, H.O., Risk Factors of Recurrent Febrile Convulsions in Children in AL-batool Teaching Hospital. *Indian Journal of Forensic Medicine & Toxicology*, 2021. 15(3).
17. Ma, X., et al., Clinical characteristics and follow-up of benign convulsions with mild gastroenteritis among children. *Medicine*, 2019. 98(2).
18. Ahn, S.H., et al., Postvaccination fever response rates in children derived using the fever coach mobile app: a retrospective observational study. *JMIR mHealth and uHealth*, 2019. 7(4): p. e12223.
19. Alruwaili, R.M.R., et al., Convulsions in Children, Arar, Northern Saudi Arabia. *The Egyptian Journal of Hospital Medicine*, 2018. 71(3): p. 2738-2741.
20. Alqudah, M.A., et al., Understanding presentations of children with fever to a Sydney emergency department. *Journal of Nursing Research and Practice*, 2019.
21. Wang, D., et al., Prognostic factors for the recurrence of afebrile seizures after benign convulsions associated with mild gastroenteritis. *Epilepsia*, 2021. 62(12): p. 3068-3075.
22. Prakash, K. and C.K. Jain, Utility of Knerr's Repertory in the Management of Febrile Convulsions. *Journal of Medical and Pharmaceutical Innovation*, 2021. 8(41): p. 5-8.
23. Baran, G. and E. Turan, Investigation of the effect of the training on fever and febrile convulsion management given to pediatric nurses on their knowledge level. *International journal of caring sciences*, 2018. 11(1): p. 478-487.
24. Laino, D., E. Mencaroni, and S. Esposito, Management of pediatric febrile seizures. *International journal of environmental research and public health*, 2018. 15(10): p. 2232.
25. Mosili, P., et al., The pathogenesis of fever-induced febrile seizures and its current state. *Neuroscience Insights*, 2020. 15: p. 2633105520956973.
26. Abd El-Moneim, H.M., M.B.E. Amir, and W. Salah, Copeptin as a Serum Biomarker for Febrile Convulsions.
27. QIAN, Q., et al., Fever, unilateral convulsion status, disturbance of consciousness. *Chinese Journal of Applied Clinical Pediatrics*, 2019: p. 781-784.
28. Subbarao, P., et al., To study zinc deficiency as a risk factor for febrile convulsions. *Journal of Evolution of Medical and Dental Sciences*, 2019. 8(15): p. 1208-1212.
29. Hameed, Z.A.A., et al., Relation of iron and zinc deficiencies to the occurrence of febrile convulsions. *Journal of pediatric neurosciences*, 2019. 14(2): p. 61.
30. Hawas, A.F., H.H. Al-Shalah, and A.H. Al-Jothary, The impact of electrolytes in pathogenesis of simple febrile convulsions. *Medical Journal of Babylon*, 2018. 15(1): p. 12.
31. Sudhakar, B. and V. Akkala, A study on febrile convulsions with special reference to incidence of bacteremia. *Cough*. 21: p. 42.
32. Schäfer, W., T. Reinders, and T. Schink, Second dose of measles-mumps-rubella-varicella vaccine (MMRV) and the risk of febrile convulsions. *Vaccine*, 2022. 40(14): p. 2168-2172.
33. Lee, H.G., et al., Febrile convulsions during recovery after anesthesia in an infant with history of MMR vaccination: A case report. *Medicine*, 2019. 98(35).
34. Pineda Solas, V., Fever and childhood vaccination. *Vacunas (English Edition)*, 2020. 21(2): p. 105-110.